Amendments to the Specification

Please amend the specification at page 2, lines 18-26, by deleting the section entitled "BRIEF DESCRIPTION OF THE DRAWINGS" as follows:

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 represents a standard curve for the screening assay using benzoylecgonine calibrators.

Fig. 2 represents a standard curve for the screening assay using morphine calibrators.

Fig. 3 represents a standard curve for the screening assay as relative concentration.

Please amend the paragraph on page 31, lines 1-2, as follows:

Pierce, Rockford, IL

TWEEN Tween-20® surfactant (#28320)

Please amend the paragraph on page 48, lines 26-29, as follows:

1. Beads were prepared in 1.5 ml microfuge tubes. Reagents were added in the order and amounts as follows: 55 μl water, 50 μl 1 M MES (pH 6.0), 5 μl 10% TWEEN Tween-20® surfactant, 100 μl beads (100 mg/ml), 280 μl anti-digoxin and 10 μl NaBH₃CN. Samples were vortexed after each addition.

Please amend the paragraph on page 53, lines 3-8, as follows:

A standard curve for the screening assay using benzoylecgonine calibrators is shown in Fig. 1. A standard curve for the screening assay using morphine calibrators in Fig. 2. A standard curve for the screening assay as relative concentration in Fig. 3. The above results demonstrate that a screening assay in accordance with the present invention for morphine combined with cocaine gives accurate, reliable results.